

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12609 A1

(51) International Patent Classification: C07D 231/54,
401/04, 413/04, 401/12, 409/12, A61K 31/415

(21) International Application Number: PCT/US00/22727

(22) International Filing Date: 19 August 2000 (19.08.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/378,234 19 August 1999 (19.08.1999) US
Not furnished 18 August 2000 (18.08.2000) US

(71) Applicant (for all designated States except US): SIGNAL
PHARMACEUTICALS, INC. [US/US]; 5555 Oberlin
Drive, San Diego, CA 92121 (US).

(72) Inventors; and


(75) Inventors/Applicants (for US only): BENNETT, Bry-
don, L. [NZ/US]; 4931 Diane Court, San Diego, CA 92117
(US). BHAGWAT, Shripad, S. [US/US]; 5015 Ashley
Falls Court, San Diego, CA 92130 (US). MANNING,
Anthony, M. [NZ/US]; 12333 Goldfish Court, San Diego,
CA 92129 (US). MURRAY, Brian, W. [US/US]; 2685
Narcissus Drive, San Diego, CA 92106 (US). O'LEARY,
Eoin, C. [IE/US]; 4964 Vista Place, San Diego, CA 92116
(US). SATOH, Yoshitaka [JP/US]; 11460 Larmier Circle,
San Diego, CA 92131 (US).

(74) Agents: HERMANN, Karl, R. et al.; Seed Intellectual
Property Law Group PLLC, Suite 6300, 701 Fifth Avenue,
Seattle, WA 98104-7092 (US).


(81) Designated States (national): AE, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,

[Continued on next page]

(54) Title: PYRAZOLOANTHRONE AND DERIVATIVES THEREOF AS JNK INHIBITORS AND THEIR COMPOSITIONS



(57) Abstract: Compounds having activity as selec-
tive inhibitors of JNK are disclosed. The compounds
of this invention are pyrazoloanthrone and derivatives
thereof having structure (I) wherein R₁ and R₂ are as
defined herein. Such compounds have utility in the
treatment of a wide range of conditions that are re-
sponsive to JNK inhibition. Thus, methods of treat-
ing such conditions are also disclosed, as are pharma-
ceutical compositions containing one or more com-
pounds of the above compounds.



WO 01/12609 A1

ATTORNEY DOCKET NUMBER: 9916-000-999

(Cam No. 501872-999079)

SERIAL NUMBER: 10/693,793

REFERENCE: B09

IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

— With international search report.

- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

PYRAZOLOANTHRONE AND DERIVATIVES THEREOF AS JNK INHIBITORS AND THEIR COMPOSITIONS

TECHNICAL FIELD

5 This invention is generally directed to pyrazoloanthrone and derivatives thereof which have utility over a wide range of indications, including activity as Jun N-terminal kinase inhibitors, and related compositions and methods.

BACKGROUND OF THE INVENTION

10 The Jun N-terminal kinase (JNK) pathway is activated by exposure of cells to environmental stress or by treatment of cells with pro-inflammatory cytokines. Targets of the JNK pathway include the transcription factors c-jun and ATF2 (Whitmarsh A.J., and Davis R.J. *J. Mol. Med.* 74:589-607, 1996). These transcription factors are members of the basic leucine zipper (bZIP) group that bind as homo- and
15 hetero-dimeric complexes to AP-1 and AP-1-like sites in the promoters of many genes (Karin M., Liu Z.G. and Zandi E. *Curr Opin Cell Biol* 9:240-246, 1997). JNK binds to the N-terminal region of c-jun and ATF-2 and phosphorylates two sites within the activation domain of each transcription factor (Hibi M., Lin A., Smeal T., Minden A., Karin M. *Genes Dev.* 7:2135-2148, 1993; Mohit A.A., Martin M.H., and Miller C.A.
20 *Neuron* 14:67-75, 1995)]. Three JNK enzymes have been identified as products of distinct genes (Hibi et al, *supra*; Mohit et al., *supra*). Ten different isoforms of JNK have been identified. These represent alternatively spliced forms of three different genes: JNK1, JNK2 and JNK3. JNK1 and 2 are ubiquitously expressed in human tissues, whereas JNK3 is selectively expressed in the brain, heart and testis (Dong, C.,
25 Yang, D., Wysk, M., Whitmarsh, A., Davis, R., Flavell, R. *Science* 270:1-4, 1998). Gene transcripts are alternatively spliced to produce four-JNK1 isoforms, four-JNK2 isoforms and two-JNK3 isoforms. JNK1 and 2 are expressed widely in mammalian tissues, whereas JNK3 is expressed almost exclusively in the brain. Selectivity of JNK signaling is achieved via specific interactions of JNK pathway components and by use
30 of scaffold proteins that selectively bind multiple components of the signaling cascade.

JIP-1 (JNK-interacting protein-1) selectively binds the MAPK module, MLK → JNKK1 → JNK.^{12,13} It has no binding affinity for a variety of other MAPK cascade enzymes. Different scaffold proteins are likely to exist for other MAPK signaling cascades to preserve substrate specificity.

- 5 JNKs are activated by dual phosphorylation on Thr-183 and Tyr-185. JNKK1 (also known as MKK 4) and JNKK2 (MKK7), two MAPKK level enzymes, can mediate JNK activation in cells (Lin A., Minden A., Martinetto H., Claret F.-Z., Lange-Carter C., Mercurio F., Johnson G.L., and Karin M. *Science* 268:286-289, 1995; Tournier C., Whitmarsh A.J., Cavanagh J., Barrett T., and Davis R.J. *Proc. Nat. Acad.*
10 *Sci. USA* 94:7337-7342, 1997). JNKK2 specifically phosphorylates JNK, whereas JNKK1 can also phosphorylate and activate p38. Both JNKK1 and JNKK2 are widely expressed in mammalian tissues. JNKK1 and JNKK2 are activated by the MAPKKK enzymes, MEKK1 and 2 (Lange-Carter C.A., Pleiman C.M., Gardner A.M., Blumer K.J., and Johnson G.L. *Science* 260:315-319, 1993; Yan M., Dai J.C., Deak J.C.,
15 Kyriakis J.M., Zon L.I., Woodgett J.R., and Templeton D.J. *Nature* 372:798-781, 1994). Both MEKK1 and MEKK2 are widely expressed in mammalian tissues.

- Activation of the JNK pathway has been documented in a number of disease settings, providing the rationale for targeting this pathway for drug discovery. In addition, molecular genetic approaches have validated the pathogenic role of this
20 pathway in several diseases. For example, autoimmune and inflammatory diseases arise from the over-activation of the immune system. Activated immune cells express many genes encoding inflammatory molecules, including cytokines, growth factors, cell surface receptors, cell adhesion molecules and degradative enzymes. Many of these genes are regulated by the JNK pathway, through activation of the transcription factors
25 AP-1 and ATF-2, including TNF α , IL-2, E-selectin and matrix metalloproteinases such as collagenase-1 (Manning A.M. and Mercurio F. *Exp Opin Invest Drugs* 6: 555-567, 1997). Monocytes, tissue macrophages and tissue mast cells are key sources of TNF α production. The JNK pathway regulates TNF α production in bacterial lipopolysaccharide-stimulated macrophages, and in mast cells stimulated through the
30 Fc ϵ RII receptor (Swanetek J.L., Cobb M.H., Geppert T.D. *Mol. Cell. Biol.* 17:6274-

6282, 1997; Ishizuka, T., Tereda N., Gerwins, P., Hamelmann E., Oshiba A., Fanger G.R., Johnson G.L., and Gelfand E.W. *Proc. Nat. Acad. Sci. USA* 94:6358-6363, 1997). Inhibition of JNK activation effectively modulates TNF α secretion from these cells. The JNK pathway therefore regulates production of this key pro-inflammatory cytokine. Matrix metalloproteinases (MMPs) promote cartilage and bone erosion in rheumatoid arthritis, and generalized tissue destruction in other autoimmune diseases. Inducible expression of MMPs, including MMP-3 and MMP-9, type II and IV collagenases, are regulated via activation of the JNK pathway and AP-1 (Gum, R., Wang, H., Lengyel, E., Juarez, J., and Boyd, D. *Oncogene* 14:1481-1493, 1997). In human rheumatoid synoviocytes activated with TNF α , IL-1, or Fas ligand the JNK pathway is activated (Han Z., Boyle D.L., Aupperle K.R., Bennett B., Manning A.M., Firestein G.S. *J. Pharm. Exp. Therap.* 291:1-7, 1999; Okamoto K., Fujisawa K., Hasunuma T., Kobata T., Sumida T., and Nishioka K. *Arth & Rheum* 40: 919-926, 1997). Inhibition of JNK activation results in decreased AP-1 activation and collagenase-1 expression (Han et al., *supra*). The JNK pathway therefore regulates MMP expression in cells involved in rheumatoid arthritis.

Inappropriate activation of T lymphocytes initiates and perpetuates many autoimmune diseases, including asthma, inflammatory bowel disease and multiple sclerosis. The JNK pathway is activated in T cells by antigen stimulation and CD28 receptor co-stimulation and regulates production of the growth factor IL-2 and cellular proliferation (Su B., Jacinto E., Hibi M., Kallunki T., Karin M., Ben-Neriah Y. *Cell* 77:727-736, 1994; Faris M., Kokot N., Lee L., and Nel A.E. *J. Biol. Chem.* 271:27366-27373, 1996). Peripheral T cells from mice genetically deficient in JNKK1 show decreased proliferation and IL-2 production after CD28 co-stimulation and PMA / Ca²⁺ ionophore activation, providing important validation for the role of the JNK pathway in these cells (Nishina H., Bachmann M., Oliveria-dos-Santos A.J., et al. *J. Exp. Med.* 186: 941-953, 1997). It is known that T cells activated by antigen receptor stimulation in the absence of accessory cell-derived co-stimulatory signals lose the capacity to synthesize IL-2, a state called clonal anergy. This is an important process by which auto-reactive T cell populations are eliminated from the peripheral circulation. Of note, anergic T cells

fail to activate the JNK pathway in response to CD3- and CD28-receptor co-stimulation, even though expression of the JNK enzymes is unchanged (Li W., Whaley C.D., Mondino A., and Mueller D.L. *Science* 271: 1272-1276, 1996). Recently, the examination of JNK-deficient mice revealed that the JNK pathway plays a key role in T cell activation and differentiation to T helper 1 and 2 cell types. JNK 1 or JNK2 knockout mice develop normally and are phenotypically unremarkable. Activated naïve CD4+ T cells from these mice fail to produce IL-2 and do not proliferate well (Sabapathy, K, Hu, Y, Kallunki, T, Schreiber, M, David, J-P, Jochum, W, Wagner, E, Karin, M. *Curr Biol* 9: 116-125, 1999). It is possible to induce T cell differentiation in T cells from these mice, generating Th1 cells (producers of IFN- γ and TNF β) and Th2 effector cells (producers of IL-4, IL-5, IL-6, IL-10 and IL-13) [22,23]. Deletion of either JNK1 or JNK2 in mice resulted in a selective defect in the ability of Th1 effector cells to express IFN γ . This suggests that JNK1 and JNK2 do not have redundant functions in T cells and that they play different roles in the control of cell growth, differentiation and death. The JNK pathway therefore, is an important point for regulation of T cell responses to antigen.

Cardiovascular disease (CVD) accounts for nearly one quarter of total annual deaths worldwide. Vascular disorders such as atherosclerosis and restenosis result from dysregulated growth of the vessel wall, restricting blood flow to vital organs. The JNK pathway is activated by atherogenic stimuli and regulates local cytokine and growth factor production in vascular cells (Yang, DD, Conze, D, Whitmarsh, AJ, et al, *Immunity*, 9:575, 1998). In addition, alterations in blood flow, hemodynamic forces and blood volume lead to JNK activation in vascular endothelium, leading to AP-1 activation and pro-atherosclerotic gene expression (Aspenstrom P., Lindberg U., and Hall A. *Curr. Biol.* 6:70-77, 1996). Ischemia and ischemia coupled with reperfusion in the heart, kidney or brain results in cell death and scar formation, which can ultimately lead to congestive heart failure, renal failure or cerebral dysfunction. In organ transplantation, reperfusion of previously ischemic donor organs results in acute leukocyte-mediated tissue injury and delay of graft function. The JNK pathway is activated by ischemia and reperfusion (Li Y., Shyy J., Li S., Lee J., Su B.,

Karin M., Chien S *Mol. Cell. Biol.* 16:5947-5954, 1996), leading to the activation of JNK-responsive genes and leukocyte-mediated tissue damage. In a number of different settings JNK activation can be either pro- or anti-apoptotic. JNK activation is correlated with enhanced apoptosis in cardiac tissues following ischemia and
5 reperfusion (Pombo CM, Bonventre JV, Avruch J, Woodgett JR, Kyriakis J.M, Force T. *J. Biol. Chem.* 269:26546-26551, 1994).

Cancer is characterized by uncontrolled growth, proliferation and migration of cells. Cancer is the second leading cause of death with 500,000 deaths and an estimated 1.3 million new cases in the United States in 1996. The role of signal
10 transduction pathways contributing to cell transformation and cancer is a generally accepted concept. The JNK pathway leading to AP-1 appears to play a critical role in cancer. Expression of c-jun is altered in early lung cancer and may mediate growth factor signaling in non-small cell lung cancer (Yin T., Sandhu G., Wolfgang C.D., Burrier A., Webb R.L., Rigel D.F. Hai T., and Whelan J. *J. Biol. Chem.* 272:19943-
15 19950, 1997). Indeed, over-expression of c-jun in cells results in transformation, and blocking c-jun activity inhibits MCF-7 colony formation (Szabo E., Riffe M., Steinberg S.M., Birrer M.J., Linnoila R.I. *Cancer Res.* 56:305-315, 1996). DNA-damaging agents, ionizing radiation and tumor necrosis factor activate the JNK pathway. In addition to regulating c-jun production and activity, JNK activation can regulate
20 phosphorylation of p53, and thus can modulate cell cycle progression (Chen T.K., Smith L.M., Gebhardt D.K., Birrer M.J., Brown P.H. *Mol. Carcinogenesis* 15:215-226, 1996). The oncogene BCR-Abl, associated with t(9,22) Philadelphia chromosome translocation of chronic myelogenous leukemia, activates JNK and leads to transformation of hematopoietic cells (Milne D.M., Campbell L.E., Campbell D.G.,
25 Meek D.W. *J. Biol. Chem.* 270:5511-5518, 1995). Selective inhibition of JNK activation by a naturally occurring JNK inhibitory protein, called JIP-1, blocks cellular transformation caused by BCR-Abl expression (Raitano A.B., Halpern J.R., Hambuch T.M., Sawyers C.L. *Proc. Nat. Acad. Sci USA* 92:11746-11750, 1995). Thus, JNK inhibitors may block transformation and tumor cell growth.


30 Accordingly, there is a need in the art for selective inhibitors of JNK, as well as for methods for preparation thereof, pharmaceutical compositions comprising

such inhibitors, and methods of inhibiting JNK's and treating diseases in mammals which are responsive to JNK inhibition. The present invention fulfills these needs, and provides further related advantages.

5 SUMMARY OF THE INVENTION

In brief, the present invention is directed to compounds having activity as selective inhibitors of JNK, as well as to compositions and methods related thereto. The compounds of the present invention (also referred to herein as "JNK inhibitors") may generally be classified as "pyrazoloanthrone derivatives" having the following

10 structure (I):



(I)

15 wherein R_1 and R_2 are as defined below, including pharmaceutically acceptable salts thereof.

The present invention is also directed to methods for treating a variety of conditions by administering an effective amount of a JNK inhibitor to an animal or subject in need thereof (referred to herein as a "patient"), typically a warm-blooded
20 animal (including a human). Prior to administration, the compounds of this invention are preferably formulated as a pharmaceutical composition which contains an effective dosage amount of one or more JNK inhibitors in combination with one (or more) pharmaceutically acceptable carrier(s). Conditions that may be treated by the compounds of this invention, or a pharmaceutical composition containing the same,
25 include any condition which may benefit from administration of JNK inhibitors, and are particularly useful for the prevention and/or treatment of various diseases including (but not limited to) rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gout, asthma,

bronchitis, cystic fibrosis, inflammatory bowel disease, irritable bowel syndrome, mucous colitis, ulcerative colitis, Crohn's disease, gastritis, esophagitis, hepatitis, multiple sclerosis, atherosclerosis, restenosis following angioplasty, left ventricular hypertrophy, myocardial infarction, stroke, ischemic damages to heart, kidney, liver, and brain, transplant rejection, endotoxin shock, psoriasis, eczema, dermatitis, epilepsy, Alzheimer's disease, Huntington's disease, Amyotrophic lateral sclerosis, peripheral neuropathies, spinal cord damage, Parkinson's disease, and cancer.

These and other aspects of this invention will be apparent upon reference to the following detailed description. To that end, certain patent and other documents are cited herein to more specifically set forth various aspects of this invention. Each of these documents are hereby incorporated by reference in their entirety.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 illustrates the ability of a representative compound of this invention to inhibit IL-2 in Jurkat T-Cell.

Figure 2 illustrates the ability of a representative compound of this invention to inhibit TNF- α in a mouse model of endotoxin shock.

Figure 3 illustrates the ability of a representative compound of this invention to inhibit leukocyte recruitment in rat model for inflamed lung.


Figure 4 illustrates the ability of a representative compound of this invention to inhibit paw swelling (Figure 4A), joint destruction (Figure 4B), transcription factor AP-1 activation (Figure 4C), and expression of MMP-13 (Figure 4D) in a rat model for adjuvant arthritis.

Figure 5 illustrates the ability of a representative compound of this invention to reduce kainic acid-induced seizure response.

DETAILED DESCRIPTION OF THE INVENTION


As mentioned above, the present invention is directed to compounds which have activity as selective inhibitors of JNK, as well as to compositions and

methods relating to the same. The compounds of this invention have the following structure (I):



and pharmaceutically acceptable salts thereof, wherein:

R_1 and R_2 are optional substituents that are the same or different and independently represent alkyl, halogen, nitro, trifluoromethyl, sulfonyl, carboxyl, alkoxy, aryl, aryloxy, arylalkyloxy, arylalkyl, cycloalkylalkyloxy, cycloalkyloxy, alkoxyalkyl, alkoxyalkoxy, aminoalkoxy, mono- or di-alkylaminoalkoxy, or a group represented by formula (a), (b), (c) or (d):



R_3 and R_4 taken together represent alkylidene or a heteroatom-containing alkylidene, or R_3 and R_4 are the same or different and independently represent hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, aryloxyalkyl, alkoxyalkyl, alkoxyamino, or alkoxy(mono- or di-alkylamino); and

R_5 represents hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, alkoxy, amino, mono- or di-alkylamino, arylamino, arylalkylamino, cycloalkylamino, or cycloalkylalkylamino.

As used herein, the terms used above having following meaning.

"Alkyl" means a straight chain or branched, saturated or unsaturated alkyl chain having from 1 to 8 carbon atoms, such as methyl, ethyl, n-propyl, iso-
5 propyl, n-butyl, iso-butyl, tert-butyl, propylenyl, 1-butenyl, propynyl, and the like.

"Halogen" means fluorine, chlorine, bromine or iodine.

"Trifluoromethyl" means $-\text{CF}_3$.

"Sulfonyl" means $-\text{SO}_2\text{H}$;

"Carboxyl" means $-\text{COOH}$.

10 "Alkoxy" means $-\text{O}-(\text{alkyl})$, such as methoxy, ethoxy, n-propyloxy, iso-propyloxy, n-butyloxy, iso-butyloxy, and the like.

"Alkoxyalkoxy" means $-\text{O}-(\text{alkyl})-\text{O}-(\text{alkyl})$, such as $-\text{OCH}_2\text{CH}_2\text{OCH}_3$, and the like.

"Alkoxy carbonyl" means $-\text{C}(=\text{O})\text{O}-(\text{alkyl})$, such as $-\text{C}(=\text{O})\text{OCH}_3$,
15 $-\text{C}(=\text{O})\text{OCH}_2\text{CH}_3$, and the like.

"Alkoxyalkyl" means $-(\text{alkyl})-\text{O}-(\text{alkyl})$, such as $-\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{OCH}_2\text{CH}_3$, and the like.

"Aryl" means a carbocyclic or heterocyclic aromatic group containing from 5 to 10 ring atoms. The ring atoms of a carbocyclic aryl group are all carbon
20 atoms, and includes phenyl and naphthyl. The ring atoms of a heterocyclic aryl group contains at least one heteroatom selected from nitrogen, oxygen and sulfur, and include pyridinyl, pyrimidinyl, furanyl, thienyl, imidazolyl, thiazolyl, pyrazolyl, pyridazinyl, pyrazinyl, triazinyl, tetrazolyl, and indolyl.

"Aryloxy" means $-\text{O}-(\text{aryl})$, such as $-\text{O}-\text{phenyl}$, $-\text{O}-\text{pyridinyl}$, and the
25 like.

"Arylalkyl" means $-(\text{alkyl})-(\text{aryl})$, such as benzyl (*i.e.*, $-\text{CH}_2\text{phenyl}$), $-\text{CH}_2\text{-pyridinyl}$, and the like.

"Arylalkyloxy" means $-\text{O}-(\text{alkyl})-(\text{aryl})$, such as $-\text{O}-\text{benzyl}$, $-\text{O}-\text{CH}_2\text{-pyridinyl}$, and the like.

"Cycloalkyl" means a cyclic alkyl having from 3 to 7 carbon atoms, such as cyclopropyl, cyclopentyl, cyclohexyl, and the like.

"Cycloalkyloxy" means -O-(cycloalkyl), such as -O-cyclohexyl, and the like.

5 "Cycloalkylalkyloxy" means -O-(alkyl)-(cycloalkyl), such as -OCH₂cyclohexyl, and the like.

"Alkylidene" means the divalent radical -C_nH_{2n}-, wherein n is an integer from 1 to 8, such as -CH₂-, -CH₂CH₂-, -CH₂-CH₂-CH₂-, -CH₂CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂CH₂-, and the like.

10 "Heteroatom-containing alkylidene" means an alkylidene wherein at least one carbon atom is replaced by a heteroatom selected from nitrogen, oxygen or sulfur, such as -CH₂CH₂OCH₂CH₂-, and the like.

"Aminoalkoxy" means -O-(alkyl)-NH₂, such as -OCH₂NH₂, -OCH₂CH₂NH₂, and the like.

15 "Mono- or di-alkylamino" means -NH(alkyl) or -N(alkyl)(alkyl), respectively, such as -NHCH₃, -N(CH₃)₂, and the like.

"Mono- or di-alkylaminoalkoxy" means -O-(alkyl)-NH(alkyl) or -O-(alkyl)-N(alkyl)(alkyl), respectively, such as -OCH₂NHCH₃, -OCH₂CH₂N(CH₃)₂, and the like.

20 "Arylamino" means -NH(aryl), such as -NH-phenyl, -NH-pyridinyl, and the like.

"Arylalkylamino" means -NH-(alkyl)-(aryl), such as -NH-benzyl, -NHCH₂-pyridinyl, and the like.

25 "Alkylamino" means -NH(alkyl), such as -NHCH₃, -NHCH₂CH₃, and the like.

"Cycloalkylamino" means -NH-(cycloalkyl), such as -NH-cyclohexyl, and the like.

"Cycloalkylalkylamino" means -NH-(alkyl)-(cycloalkyl), such as -NHCH₂-cyclohexyl, and the like.


In the embodiment wherein R_1 and R_2 are not present, compounds of this invention have the following structure (II) (also referred to herein as "Compound 1"):




(II)

This compound is commercially available from Pfaltz-Bauer (Conn., U.S.).

In the embodiment wherein only one of R_1 and R_2 is present, compounds of this invention have one of the following structures (III) or (IV):




(III)




(IV)


In the embodiment wherein both R_1 and R_2 are present, compounds of this invention have one of the following structures (V), (VI) or (VII):



(V)



(VI)




(VII)

Pharmaceutically acceptable salts of compounds of structure (I) are also within the scope of this invention. To this end, the compound may generally be utilized as the free base. Alternatively, the compounds may be used in the form of acid addition salts. Acid addition salts of the free base amino compounds of the present invention
5 may be prepared by methods well known in the art, and may be formed from organic and inorganic acids. Suitable organic acids include maleic, fumaric, benzoic, ascorbic, succinic, methanesulfonic, acetic, oxalic, propionic, tartaric, salicylic, citric, gluconic, lactic, mandelic, cinnamic, aspartic, stearic, palmitic, glycolic, glutamic, and benzenesulfonic acids. Suitable inorganic acids include hydrochloric, hydrobromic,
10 sulfuric, phosphoric, and nitric acids. Thus, the term "pharmaceutically acceptable salt" of a compound of structure (I) is intended to encompass any and all acceptable salt forms.

The compounds of this invention may generally be made by organic synthesis techniques known to those skilled in the art, as well as by the following
15 general techniques and by the procedures set forth in the Examples. To that end, the compounds of this invention may be made according to the following Reaction Schemes 1 through 7.

Reaction Scheme 1

20




In Reaction Scheme 1, pyrazoloanthrones of this invention may be prepared by condensation of appropriate anthraquinones having a leaving group at the
25 1-position (such as fluoro, chloro, bromo, iodo, nitro, methanesulfonyloxy, tosyloxy or phenoxy) with hydrazine in a suitable solvent (such as pyridine, dimethylformamide, methylene chloride, chloroform, or dioxane). The reaction is carried out at temperatures


ranging 0°C to 200°C for 1 to 16 hours. Suitable anthraquinone starting materials are commercially available from a variety of sources with the R₁ and/or R₂ groups at various positions on the anthraquinone ring. For purpose of illustration, the following reaction schemes depict synthesis of 5- and/or 7-substituted pyrazoloanthrones. One skilled in the art will recognize that pyrazoloanthrones substituted at other positions may be made in a similar manner from the appropriately substituted pyrazoloanthrone starting material.

Reaction Scheme 2


10



In Reaction Scheme 2, pyrazoloanthrones with 5-amino substituents may be prepared by condensation of 5-chloropyrazoloanthrone with mono- or disubstituted amines at 0 to 250°C for 1 to 16 hours, either in the absence or the presence of a solvent. Typically solvents are pyridine, dimethylformamide, dimethylsulfoxide, dichloroethane, chloroform, tetrahydrofuran, dioxane, diglyme, or triglyme in the presence of excess amount of the amine, or in the presence of an acid quenching agent such as triethylamine, diisopropylethylamine, sodium bicarbonate, potassium carbonate, or sodium hydroxide.

Reaction Scheme 3


- 5 In Reaction Scheme 3, pyrazoloanthrones with 7-amino substituents may be prepared by condensation of 7-chloropyrazoloanthrone with mono- or disubstituted amines at 0 to 250°C for 1 to 16 hours either in the absence or the presence of a solvent. Typically solvents are pyridine, dimethylformamide, dimethylsulfoxide, dichloroethane, chloroform, tetrahydrofuran, dioxane, diglyme, or triglyme in the presence of excess
- 10 amount of the amine, or in the presence of an acid quenching agent such as triethylamine, diisopropylethylamine, sodium bicarbonate, potassium carbonate, or sodium hydroxide.

Reaction Scheme 4

- 5 In Reaction Scheme 4, pyrazoloanthrones with 5-acyl- or sulfonylamino substituents may be prepared by condensation of 5-amino-2-(2-methoxyethoxymethyl)pyrazoloanthrone with acid chlorides and sulfonyl chlorides followed by the deprotection. Condensation of 5-amino-2-(2-methoxyethoxymethyl)pyrazoloanthrone with appropriate acid chlorides R_6COCl or
- 10 sulfonyl chlorides $\text{R}_6\text{SO}_2\text{Cl}$ is carried out in the presence of an acid quenching agent such as triethylamine, diisopropylethylamine, sodium bicarbonate, potassium carbonate, or sodium hydroxide at -20 to 50°C for 0.5 to 16 hours in solvents such as methylene chloride, chloroform, tetrahydrofuran, dioxane, dimethylformamide, and ethyl acetate. The deprotection step may be performed by the treatment of the product mentioned
- 15 above with an acid such as trifluoroacetic acid, aqueous hydrochloric acid, aqueous hydrobromic acid, or aqueous sulfuric acid.

The starting material may be prepared in two steps. The 2-position of 5-nitropyrzoloanthrone may be protected by a protective group such as methoxymethyl (MOM), methoxyethoxymethyl (MEM), 2-trimethylsilylethoxymethyl (SEM), or 4-methoxybenzyl (PMB) with an aid of a base such as triethylamine, diisopropylethylamine, pyridine, sodium hexamethyldisilazide, potassium hexamethyldisilazide, or lithium diisopropylamide. 4-(N, N-dimethylamino)pyridine (DMAP) may be used as a catalyst when a tertiary amine is used as a base. The reaction is typically carried out at -40 to 60°C for 1 to 16 hours in a solvent such as methylene chloride, chloroform, tetrahydrofuran, dioxane, or dimethoxyethane. As the nitrogen protective group, MEM group is preferred.


N-Protected 5-nitropyrzoloanthrone is then reduced to its 5-amino derivative by a variety of reducing agents such as Sn or Fe metal in acidic media such as acetic acid or aqueous hydrochloric acid. The reaction is typically run at 20 to 160°C for 1 to 16 hours. The same transformation can be carried out by hydrogenation in the presence of a transition-metal catalyst such as palladium, platinum, rhodium, or iridium with or without a support such as charcoal in a solvent such as ethanol, ethyl acetate, tetrahydrofuran, dioxane, or dimethoxyethane at 1 to 20 atmospheres of hydrogen at 20 to 60°C for 1 to 16 hours. The procedure using hydrogenation with palladium on charcoal as the catalyst is preferred.

Reaction Scheme 5

- 5 In Reaction Scheme 5, pyrazoloanthrones with 7-acyl- or sulfonylamino substituents may be prepared by condensation of 7-amino-2-(2-methoxyethoxymethyl)pyrazoloanthrone with acid chlorides and sulfonyl chlorides followed by the deprotection. Condensation of 7-amino-2-(2-methoxyethoxymethyl)pyrazoloanthrone with appropriate acid chlorides R_6COCl or
- 10 sulfonyl chlorides R_6SO_2Cl is carried out in the presence of an acid quenching agent such as triethylamine, diisopropylethylamine, sodium bicarbonate, potassium carbonate, or sodium hydroxide at -20 to $50^\circ C$ for 0.5 to 16 hours in solvents such as methylene chloride, chloroform, tetrahydrofuran, dioxane, dimethylformamide, or ethyl acetate. The deprotection step may be performed by the treatment of the product mentioned
- 15 above with an acid such as trifluoroacetic acid, aqueous hydrochloric acid, aqueous hydrobromic acid, or aqueous sulfuric acid.

The starting material is prepared in two steps. The 2-position of 7-nitropyrzoloanthrone is protected by a protective group such as methoxymethyl (MOM), methoxyethoxymethyl (MEM), 2-trimethylsilylethoxymethyl (SEM), or 4-methoxybenzyl (PMB) with an aid of a base such as triethylamine, diisopropylethylamine, pyridine, sodium hexamethyldisilazide, potassium hexamethyldisilazide, or lithium diisopropylamide. 4-(N, N-dimethylamino)pyridine (DMAP) can be used as a catalyst when a tertiary amine is used as a base. The reaction is typically carried out at -40 to 60°C for 1 to 16 hours in a solvent such as methylene chloride, chloroform, tetrahydrofuran, dioxane, or dimethoxyethane. As the nitrogen protective group, MEM group is preferred.

N-Protected 7-nitropyrzoloanthrone is then reduced to its 7-amino derivative by a variety of reducing agents such as Sn or Fe metal in acidic media such as acetic acid or aqueous hydrochloric acid. The reaction is typically run at 20 to 160°C for 1 to 16 hours. The same transformation can be carried out by hydrogenation in the presence of a transition-metal catalyst such as palladium, platinum, rhodium, or iridium with or without a support such as charcoal in a solvent such as ethanol, ethyl acetate, tetrahydrofuran, dioxane, or dimethoxyethane at 1 to 20 atmospheres of hydrogen at 20 to 60°C for 1 to 16 hours. The procedure using hydrogenation with palladium on charcoal as the catalyst is preferred.

Reaction Scheme 6


5 In Reaction Scheme 6, pyrazoloanthrones with 5-alkoxy substituents may be prepared by condensation of 5-hydroxy-2-(2-methoxyethoxymethyl)-pyrazoloanthrone with alkyl halides and sulfonates R_7-X followed by the deprotection. As the leaving group X, chloride, bromide, iodide, methanesulfonate, tosylate, benzenesulfonate, or triflate can be used. Condensation of 5-hydroxy-2-(2-methoxyethoxymethyl)pyrazoloanthrone with appropriate alkyl halides and sulfonates is carried out in the presence of an acid quenching agent such as triethylamine, diisopropylethylamine, sodium bicarbonate, potassium carbonate, or sodium hydroxide at -20 to 50°C for 0.5 to 16 hours in solvents such as methylene chloride, chloroform, tetrahydrofuran, dioxane, dimethylformamide, or ethyl acetate. The deprotection step is performed by the treatment of the product mentioned above with an acid such as trifluoroacetic acid, aqueous hydrochloric acid, aqueous hydrobromic acid, or aqueous sulfuric acid.

The starting material is prepared in two steps. The 2-position of 5-benzyloxy-2-(2-methoxyethoxymethyl)pyrazoloanthrone is protected by a protective group such as methoxymethyl

(MOM), methoxyethoxymethyl (MEM), 2-trimethylsilylethoxymethyl (SEM), or 4-methoxybenzyl (PMB) with an aid of a base such as triethylamine, diisopropylethylamine, pyridine, sodium hexamethyldisilazide, potassium hexamethyldisilazide, or lithium diisopropylamide. 4-(N,N-dimethylamino)pyridine (DMAP) can be used as a catalyst when a tertiary amine is used as a base. The reaction is typically carried out at -40 to 60°C for 1 to 16 hours in a solvent such as methylene chloride, chloroform, tetrahydrofuran, dioxane, or dimethoxyethane. As the nitrogen protective group, MEM group is preferred.

N-Protected 5-benzyloxypyrazoloanthrone is then reduced to its 5-hydroxy derivative by hydrogenation in the presence of a transition-metal catalyst, such as palladium platinum, rhodium, or iridium with or without a support such as charcoal in a solvent such as ethanol, ethyl acetate, tetrahydrofuran, dioxane, or dimethoxyethane at 1 to 20 atmospheres of hydrogen at 20 to 60°C for 1 to 16 hours. The procedure using hydrogenation with palladium on charcoal as the catalyst is preferred.

Reaction Scheme 7



In Reaction Scheme 7, pyrazoloanthrones with 5-alkoxy substituents may be prepared by condensation of 7-hydroxy-2-(2-methoxyethoxymethyl)-pyrazoloanthrone with alkyl halides and sulfonates R₁-X followed by the deprotection. As the leaving group X, chloride, bromide, iodide, methanesulfonate, tosylate, benzenesulfonate, or triflate can be used. Condensation of 7-hydroxy-2-(2-methoxyethoxymethyl)pyrazoloanthrone with appropriate alkyl halides and sulfonates is carried out in the presence of an acid quenching agent such as triethylamine, diisopropylethylamine, sodium bicarbonate, potassium carbonate, or sodium hydroxide at -20 to 50°C for 0.5 to 16 hours in solvents such as methylene chloride, chloroform, tetrahydrofuran, dioxane, dimethylformamide, or ethyl acetate. The deprotection step is performed by the treatment of the product mentioned above with an acid such as trifluoroacetic acid, aqueous hydrochloric acid, aqueous hydrobromic acid, or aqueous sulfuric acid.

The starting material is prepared in two steps. The 2-position of 7-benzyloxypyrazoloanthrone is protected by a protective group such as methoxymethyl (MOM), methoxyethoxymethyl (MEM), 2-trimethylsilylethoxymethyl (SEM), or 4-methoxybenzyl (PMB) with an aid of a base such as triethylamine, diisopropylethylamine, pyridine, sodium hexamethyldisilazide, potassium hexamethyldisilazide, or lithium diisopropylamide. 4-(N, N-dimethylamino)pyridine (DMAP) can be used as a catalyst when a tertiary amine is used as a base. The reaction is typically carried out at -40 to 60°C for 1 to 16 hours in a solvent such as methylene chloride, chloroform, tetrahydrofuran, dioxane, or dimethoxyethane. As the nitrogen protective group, MEM group is preferred.

N-Protected 7-benzyloxypyrazoloanthrone is then reduced to its 7-hydroxy derivative by hydrogenation in the presence of a transition-metal catalyst, such as palladium platinum, rhodium, or iridium with or without a support such as charcoal in a solvent such as ethanol, ethyl acetate, tetrahydrofuran, dioxane, or dimethoxyethane at 1 to 20 atmospheres of hydrogen at 20 to 60°C for 1 to 16 hours. The procedure using hydrogenation with palladium on charcoal as the catalyst is preferred.

Compounds of structures (V), (VI) and (VII) may be made by the same procedures as outlined above by utilizing starting materials having multiple reactive sites at the corresponding positions to the desired product.

In another embodiment of the invention, pharmaceutical compositions containing one or more compounds of this invention are disclosed. For purpose of administration, a compound of structure (I) is preferably formulated as a pharmaceutical composition. Pharmaceutical compositions of the present invention comprise a compound of this invention and a pharmaceutically acceptable carrier, wherein the compound is present in the composition in an amount which is effective to treat the condition of interest. Preferably, the pharmaceutical compositions of the present invention include a compound of structure (I) in an amount from 0.1 mg to 250 mg per dosage depending upon the route of administration, and more typically from 1 mg to 60 mg. Appropriate concentrations and dosages can be readily determined by one skilled in the art.

Pharmaceutically acceptable carriers are familiar to those skilled in the art. For compositions formulated as liquid solutions, acceptable carriers include saline and sterile water, and may optionally include antioxidants, buffers, bacteriostats and other common additives. The compositions can also be formulated as pills, capsules, granules, or tablets which contain, in addition to a compound of this invention, diluents, dispersing and surface active agents, binders, and lubricants. One skilled in this art may further formulate the compounds of this invention in an appropriate manner, and in accordance with accepted practices, such as those disclosed in *Remington's Pharmaceutical Sciences*, Gennaro, Ed., Mack Publishing Co., Easton, PA 1990.

In another embodiment, the present invention provides a method for treating a variety of conditions by administering an effective amount of a JNK inhibitor to a patient in need thereof. Conditions that may be treated by the compounds of this invention, or a pharmaceutical composition containing the same, include any condition which is responsive to JNK inhibition, and thereby benefit from administration of a JNK inhibitor. Representative conditions in this regard include (but not limited to) rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gout, asthma, bronchitis,

cystic fibrosis, inflammatory bowel disease, irritable bowel syndrome, mucous colitis, ulcerative colitis, Crohn's disease, gastritis, esophagitis, hepatitis, multiple sclerosis, atherosclerosis, restenosis following angioplasty, left ventricular hypertrophy, myocardial infarction, stroke, ischemic damage to the heart, kidney, liver, or brain, 5 transplant rejection (such as kidney, liver, heart, lung, and the like), endotoxin shock, psoriasis, eczema, dermatitis, epilepsy, Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, peripheral neuropathies, spinal cord damage, and cancer.

The methods of this invention include systemic administration of a 10 compound of this invention, preferably in the form of a pharmaceutical composition. As used herein, systemic administration encompasses both oral and parenteral methods of administration. For oral administration, suitable pharmaceutical compositions include powders, granules, pills, tablets, and capsules as well as liquids, syrups, suspensions, and emulsions. These compositions may also include flavorants, 15 preservatives, suspending, thickening and emulsifying agents, and other pharmaceutically acceptable additives. For parental administration, the compounds of the present invention can be prepared in aqueous injection solutions which may contain buffers, antioxidants, bacteriostats, and other additives commonly employed in such solutions.


20 The following examples are offered by way of illustration, not limitation.


EXAMPLES

EXAMPLE 1

Synthesis of Representative Compounds

25




A. Anthra[1,9cd]pyrazol-6(2H)-one ("Compound 1")


5

Anhydrous hydrazine is added to a solution of 2-chloroanthraquinone (Aldrich) in 10 mL pyridine, and the mixture heated at 100°C for 16 hours. The mixture is cooled and the solvent is evaporated in vacuo. The residue is taken in hot 6N HCl, and the solid is collected by filtration. Flash chromatography of the crude material
10 on silica gel affords anthra[1,9cd]pyrazol-6(2H)-one ("Compound 1") as yellow solids.

Due to limited solubility of Compound 1, purification of the same may be achieved by first derivatizing Compound 1 to a more soluble intermediate, such as the corresponding acetate, recrystallizing the intermediate, and then converting the intermediate to yield purified Compound 1 in good yield. More specifically, to solution
15 of the pyrazoloanthrone (9.67 g, 43.9 mmol) in acetic acid (700 mL) is added acetic anhydride (12.4 mL, 132 mmol). The solution is heated to 80°C for 5 hours and then cooled to room temperature. After 16 hours, the reaction is cooled to 0°C for 2 hours. The reaction is then filtered to give the N-acetylpyrazoloanthrone intermediate. This intermediate is recrystallized in acetic acid to give the pure intermediate (5.96 g, 52%).
20 ¹H NMR (CDCl₃) δ 10.6 (br s, 1H), 8.46 (d, 1H), 8.33 (d, 1H), 8.26 (d, 1H), 8.08 (d, 1H), 7.96-7.87 (m, 2H), 7.78 (t, 1H), 2.83 (s, 3H); ES-MS (m/z) 263 [M+1]⁺. To a solution of the pure intermediate (5.96 g, 23 mmol) in methanol (600 mL) is added ammonium hydroxide (60 mL). The reaction is stirred at room temperature for 16 hours and then filtered and dried in a vacuum oven. A second crop of crystals is
25 recovered to give a total of 4.8 g of Compound 1 at greater than 98% purity. ES-MS (m/z) 221 [M + 1]⁺.


B. 5-Chloroanthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner from 1,4-dichloroanthraquinone (commercial product).


C. 7-Chloroanthra[1,9cd]pyrazol-6(2H)-one

10


This compound may be made in the same manner from 1,5-dichloroanthraquinone (commercial product).

15 D. 5-Nitroanthra[1,9cd]pyrazol-6(2H)-one

This compound may be made from 1,4-dinitroanthraquinone (Krapcho,
20 A. P.; Avery, K. L., Jr. *J. Org. Chem.* 55, 5562-4, 1990).

E. 7-Nitroanthra[1.9cd]pyrazol-6(2H)-one


5 This compound may be made in the same manner from 1,5-dichloroanthraquinone (commercial product).

F. 5-Benzyloxvanthra[1.9cd]pyrazol-6(2H)-one

10

 This compound may be made in the same manner from 1-nitro-4-benzyloxyanthraquinone. This starting material may be prepared as follows. Benzyl bromide is added to 1-nitro-4-hydroxyanthraquinone (Aldrich) and potassium carbonate
15 in dimethylformamide, and the mixture is stirred for 16 hours. Water is added and the mixture is extracted with ethyl acetate (x2). The combined organic layer is washed sequentially with sodium bicarbonate solution, water, 1N hydrochloric acid, and brine, dried, and evaporated. The residue is chromatographed on silica gel to afford 1-nitro-4-benzyloxyanthraquinone.


20

G. 7-Benzyloxyanthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner from 1-nitro-5-benzyloxyanthraquinone, which starting material may prepared as disclosed in German Patent No. DE 2254199 to Reubke, Hohmann and Bien.


H. 5-(Acetylamino)anthra[1,9cd]pyrazol-6(2H)-one

10




 This compound may be made in the same manner from 4-acetylamino-1-chloroanthraquinone. This starting material may be prepared as follows. 4-Amino-1-chloroanthraquinone is taken in pyridine and treated with acetic anhydride. The mixture is stirred for 1 hour, and poured onto water. The solids are collected by
15 filtration, washed with water, and dried in vacuo to give 4-acetylamino-1-chloroanthraquinone as a colorless solid.


EXAMPLE 2

Synthesis of Representative Compounds

5

A. 5-(Dimethylamino)anthra[1,9cd]pyrazol-6(2H)-one


- 10 A mixture of 5-chloroanthra[1,9cd]pyrazol-6(2H)-one (Example 1-B) and dimethylamine in pyridine is heated at 100 °C for 16 hours. The mixture is cooled and evaporated. The residue is chromatographed on silica gel to give the desired compound as yellow solids.

15 B. 5-(1-Piperidiny)anthra[1,9cd]pyrazol-6(2H)-one

This compound may be made in the same manner using piperidine as the amine.

C. 5-(1-Morpholinyl)anthra[1,9cd]pyrazol-6(2H)-one


5



This compound may be made in the same manner using morpholine as the amine.


10

D. 5-(Benzylamino)anthra[1,9cd]pyrazol-6(2H)-one




15

This compound may be made in the same manner using benzylamine as the amine.

E. 5-[(4-Pyridylmethyl)amino]anthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner using 4-pyridylmethylaniline as the amine.


F. 5-[2-(1-Piperidinyl)ethylamino]anthra[1,9cd]pyrazol-6(2H)-one


10

This compound may be made in the same manner using 2-(1-piperidyl)ethylaniline as the amine.

15

EXAMPLE 3


Synthesis of Representative Compounds

A. 7-(Dimethylamino)anthra[1,9cd]pyrazol-6(2H)-one

5


A mixture of 6-chloroanthra[1,9cd]pyrazol-6(2H)-one (Example 1-C) and dimethylamine in pyridine is heated at 100°C for 16 hours. The mixture is cooled and evaporated. The residue is chromatographed on silica gel to give the desired compound as yellow solids.

10


B. 5-(1-Piperidinyl)anthra[1,9cd]pyrazol-6(2H)-one

15

This compound may be made in the same manner using piperidine as the amine.


C. 5-(1-Morpholinyl)anthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner using morpholine as the amine.

D. 5-(Benzylamino)anthra[1,9cd]pyrazol-6(2H)-one

10


This compound may be made in the same manner using benzylamine as the amine.

15 E. 5-((4-Pyridylmethyl)amino)anthra[1,9cd]pyrazol-6(2H)-one

This compound may be made in the same manner using 4-pyridylmethylaniline as the amine.

F. 5-(2-(1-Piperidinyl)ethylamino)anthra[1,9cd]pyrazol-6(2H)-one

5




This compound may be made in the same manner using 2-(1-piperidyl)ethylamine as the amine.

10


EXAMPLE 4

Synthesis of Representative Compounds



Y = C, n = 1
Y = S, n = 2


15

A. 5-(Benzoylamino)anthra[1,9cd]pyrazol-6(2H)-one


5 Benzoyl chloride is added to a solution of 2-(methoxyethoxymethyl)-5-aminoanthra[1,9cd]pyrazol-6-(2H)one and triethylamine in methylene chloride at 0°C. The mixture is stirred for 16 hours, quenched with water, and extracted with ethyl acetate (x2). The combined organic layer is washed with sodium bicarbonate solution, and brine, dried and evaporated. The crude reaction mixture is then taken in aqueous
10 6N hydrochloric acid, and heated at 80°C for 4 hours. After cooling, the mixture is extracted with ethyl acetate (x2), washed with brine, dried, and evaporated. The residue is chromatographed on silica gel to furnish the desired amide as a yellow solid.

The starting material is prepared as follows. Sodium hexamethyldisilazide is added to a cooled (0°C) solution of 5-nitroanthra[1,9cd]pyrazol-
15 6(2H)-one (Example 1-D) in tetrahydrofuran, and the mixture is stirred for 30 minutes at 0°C. MEM-chloride is added, and the mixture is stirred for 16 hours at room temperature. Water is added and the mixture is extracted with ethyl acetate (x2). The combined organic layer is washed with aqueous sodium bicarbonate solution, water, 1N hydrochloric acid, and brine, dried and evaporated. The residue is chromatographed on
20 silica gel to give 2-MEM-5-nitroanthra[1,9cd]pyrazol-6(2H)-one as an oil.

Palladium(10%) on charcoal and 2-MEM-5-nitroanthra[1,9cd]pyrazol-6(2H)-one in ethanol is placed under 1-atm of hydrogen, and the mixture was stirred for 6 hours. The catalyst is filtered off over celite, and the filtrate is evaporated to dryness to give 2-(methoxyethoxymethyl)-5-aminoanthra[1,9cd]pyrazol-6-(2H)one, which is
25 used without further purification.


B. 5-(Isonicotinylamino)anthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner using isonicotinoyl chloride as the acid chloride

C. 5-(Nicotinylamino)anthra[1,9cd]pyrazol-6(2H)-one

10


This compound may be made in the same manner using nicotinoyl chloride as the acid chloride.

15 D. 5-(2-Thiophenecarbonvlamino)anthra[1,9cd]pyrazol-6(2H)-one

This compound may be made in the same manner using 2-thiophenecarboxylic acid as the acid chloride.

E. 5-(3-Methylbutyrylamino)anthra[1,9cd]pyrazol-6(2H)-one


5



This compound may be made in the same manner using isopentanoyl chloride as the acid chloride.


10

F. 5-(3-Methanesulfonylamino)anthra[1,9cd]pyrazol-6(2H)-one



15

This compound may be made in the same manner using methanesulfonyl chloride as the sulfonyl chloride.


G. 5-(3-Benzenesulfonylamino)anthra[1,9cd]pyrazol-6(2H)-one


- 5 This compound may be made in the same manner using benzenesulfonyl chloride as the sulfonyl chloride.

EXAMPLE 5

Synthesis of Representative Compounds

10




A. 7-(Benzoylamino)anthra[1,9cd]pyrazol-6(2H)-one


5 Benzoyl chloride is added to a solution of 2-(methoxyethoxymethyl)-7-aminoanthra[1,9cd]pyrazol-6-(2H)one and triethylamine in methylene chloride at 0°C. The mixture is stirred for 16 hours, quenched with water, and extracted with ethyl acetate (x2). The combined organic layer is washed with sodium bicarbonate solution, and brine, dried and evaporated. The crude reaction mixture is then taken in aqueous
10 6N hydrochloric acid, and heated at 80°C for 4 hours. After cooling, the mixture is extracted with ethyl acetate (x2), washed with brine, dried, and evaporated. The residue is chromatographed on silica gel to furnish the desired amide as a yellow solid.

The starting material is prepared as follows. Sodium hexamethyldisilazide is added to a cooled (0°C) solution of 7-nitroanthra[1,9cd]pyrazol-
15 6(2H)-one (Example 1-E) in tetrahydrofuran, and the mixture is stirred for 30 minutes at 0°C. MEM-chloride is added, and the mixture is stirred for 16 hours at room temperature. Water is added and the mixture is extracted with ethyl acetate (x2). The combined organic layer is washed with aqueous sodium bicarbonate solution, water, 1N hydrochloric acid, and brine, dried and evaporated. The residue is chromatographed on
20 silica gel to give 2-MEM-7-nitroanthra[1,9cd]pyrazol-6(2H)-one as an oil.

Palladium(10%) on charcoal and 2-MEM-5-nitroanthra[1,9cd]pyrazol-6(2H)-one in ethanol is placed under 1-atm of hydrogen, and the mixture was stirred for 6 hours. The catalyst is filtered off over celite, and the filtrate is evaporated to dryness to give 2-(methoxyethoxymethyl)-7-aminoanthra[1,9cd]pyrazol-6-(2H)one, which is
25 used without further purification.


B. 7-(Isonicotinylamino)anthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner using isonicotinoyl chloride as the acid chloride.

C. 7-(Nicotinylamino)anthra[1,9cd]pyrazol-6(2H)-one

10


 This compound may be made in the same manner using nicotinoyl chloride as the acid chloride.

15 D. 5-(2-Thiophenecarbonylamino)anthra[1,9cd]pyrazol-6(2H)-one

This compound may be made in the same manner using 2-thiophenecarboxylic acid chloride as the acid chloride.

E. 7-(3-Methylbutyrylamino)anthra[1,9cd]pyrazol-6(2H)-one


5



This compound may be made in the same manner using isopentanoyl chloride as the acid chloride.


10

F. 7-(3-Methanesulfonylamino)anthra[1,9cd]pyrazol-6(2H)-one



15

This compound may be made in the same manner using methanesulfonyl chloride as the sulfonyl chloride.

G. 7-(3-Benzenesulfonylamino)anthra[1,9cd]pyrazol-6(2H)-one


5 This compound may be made in the same manner using benzenesulfonyl chloride as the sulfonyl chloride.

EXAMPLE 6

10

Synthesis of Representative CompoundsA. 5-(3-Methoxybutyloxy)anthra[1,9cd]pyrazol-6(2H)-one

15



Isopentyl bromide is added to a mixture of 3-(2-methoxyethoxymethyl)-5-hydroxyanthra[1,9cd]pyrazol-6(2H)-one and potassium carbonate in dimethylformamide at room temperature. After stirring the mixture for


20

sixteen hours, water is added, and the mixture was extracted with ethyl acetate (x2). The combined organic layer is washed with aqueous sodium bicarbonate, water, 1N hydrochloric acid, and brine, dried and evaporated. The residue is taken in 6N hydrochloric acid and heated at 80°C for 4 hours. After cooling, the mixture is
5 extracted with ethyl acetate (x2), and the combined organic layer is washed with brine, dried, and evaporated. The residue is purified by column chromatography to afford the title compound as yellow solid.

The starting material is prepared as follows. Sodium hexamethyldisilazide is added to a cooled (0°C) solution of 5-
10 benzyloxyanthra[1,9cd]pyrazol-6(2H)-one (Example 1-F) in tetrahydrofuran, and the mixture is stirred for 30 minutes at 0°C. MEM-chloride is added, and the mixture is stirred for 16 hours at room temperature. Water is added and the mixture is extracted with ethyl acetate (x2). The combined organic layer is washed with aqueous sodium bicarbonate solution, water, 1N hydrochloric acid, and brine, dried and evaporated. The
15 residue is chromatographed on silica gel to give 2-MEM-5-benzyloxyanthra[1,9cd]pyrazol-6(2H)-one as an oil.

Palladium(10%) on charcoal and 2-MEM-5-benzyloxyanthra[1,9cd]pyrazol-6(2H)-one in ethanol is placed under 1-atm of hydrogen, and the mixture stirred for 6 hours. The catalyst is filtered off over celite,
20 and the filtrate is evaporated to dryness to give 2-(2-methoxyethoxymethyl)-5-hydroxyanthra[1,9cd]pyrazol-6-(2H)one, which is used without further purification.


B. 5-(4-Pyridylmethoxy)anthra[1,9cd]pyrazol-6(2H)-one



This compound may be made in the same manner using chloromethyl-4-pyridine as the alkyl halide.

C. 5-(3-Pyridylmethoxy)anthra[1,9cd]pyrazol-6(2H)-one


5



This compound may be made in the same manner using chloromethyl-3-pyridine as the alkyl halide.

10


D. 5-(2-Methoxyethoxy)anthra[1,9cd]pyrazol-6(2H)-one



15

This compound may be made in the same manner using 2-methoxyethyl bromide as the alkyl halide.

E. 5-(2-Dimethylaminoethoxy)anthra[1,9cd]pyrazol-6(2H)-one




20


This compound may be made in the same manner using 2-dimethylaminoethyl chloride as the alkyl halide.

5

EXAMPLE 7

Synthesis of Representative Compounds

10

A. 7-(3-Methylbutyloxy)anthra[1,9cd]pyrazol-6(2H)-one

15

Isopentyl bromide is added to a mixture of 3-(2-methoxyethoxymethyl)-7-hydroxyanthra[1,9cd]pyrazol-6(2H)-one and potassium carbonate in dimethylformamide at room temperature. After stirring the mixture for sixteen hours, water is added, and the mixture was extracted with ethyl acetate (x2). The combined organic layer is washed with aqueous sodium bicarbonate, water, 1N hydrochloric acid, and brine, dried and evaporated. The residue is taken in 6N hydrochloric acid and heated at 80°C for 4 hours. After cooling, the mixture is extracted with ethyl acetate (x2), and the combined organic layer is washed with brine, dried, and evaporated. The residue is purified by column chromatography to afford the title compound as yellow solid.


20

The starting material is prepared as follows. Sodium hexamethyldisilazide is added to a cooled (0°C) solution of 7-benzyloxyanthra[1,9cd]pyrazol-6(2H)-one (Example 1-F) in tetrahydrofuran, and the mixture is stirred for 30 minutes at 0°C. MEM-chloride is added, and the mixture is stirred for 16 hours at room temperature. Water is added and the mixture is extracted with ethyl acetate (x2). The combined organic layer is washed with aqueous sodium bicarbonate solution, water, 1N hydrochloric acid, and brine, dried and evaporated. The residue is chromatographed on silica gel to give 2-MEM-7-benzyloxyanthra[1,9cd]pyrazol-6(2H)-one as an oil.


10 Palladium(10%) on charcoal and 2-MEM-7-benzyloxyanthra[1,9cd]pyrazol-6(2H)-one in ethanol is placed under 1-atm of hydrogen, and the mixture was stirred for 6 h. The catalyst is filtered off over celite, and the filtrate is evaporated to dryness to give 2-(2-methoxyethoxymethyl)-7-hydroxyanthra[1,9cd]pyrazol-6-(2H)one, which is used without further purification.

15


B. 7-(4-Pyridylmethoxy)anthra[1,9cd]pyrazol-6(2H)-one



20 This compound may be made in the same manner using chloromethyl-4-pyridine as the alkyl halide.


C. 7-(3-Pyridylmethoxy)anthra[1,9cd]pyrazol-6(2H)-one

5 This compound may be made in the same manner using chloromethyl-3-pyridine as the alkyl halide.

D. 7-(2-Methoxyethoxy)anthra[1,9cd]pyrazol-6(2H)-one

10

This compound may be made in the same manner using 2-methoxyethyl bromide as the alkyl halide.

15 E. 7-(2-Dimethylaminoethoxy)anthra[1,9cd]pyrazol-6(2H)-one

20 This compound may be made in the same manner using 2-dimethylaminoethyl chloride as the alkyl halide.

EXAMPLE 8

Activity of Representative Compound

5

The compounds of this invention may be assayed for their activity accordingly to the following procedures.

JNK Assay

- 10 To 10 μ L of the test compound in 20% DMSO/80% dilution buffer consisting of 20 mM HEPES (pH 7.6), 0.1 mM EDTA, 2.5 mM magnesium chloride, 0.004% Triton x100, 2 μ g/mL leupeptin, 20 mM β -glycerolphosphate, 0.1 mM sodium vanadate, and 2 mM DTT in water is added 30 μ L of 50-200 ng His6-JNK1, JNK2 or JNK3 in the same dilution buffer. The mixture is preincubated for 30 minutes at room
- 15 temperature. Sixty microliter of 10 μ g GST-c-Jun(1-79) in assay buffer consisting of 20 mM HEPES (pH 7.6), 50 mM sodium chloride, 0.1 mM EDTA, 24 mM magnesium chloride, 1 mM DTT, 25 mM PNPP, 0.05% Triton x100, 11 μ M ATP, and 0.5 μ Ci γ -32P ATP in water is added and the reaction is allowed to proceed for 1 hour at room temperature. The c-Jun phosphorylation is terminated by addition of 150 μ L of 12.5%
- 20 trichloroacetic acid. After 30 minutes, the precipitate is harvested onto a filter plate, diluted with 50 μ L of the scintillation fluid and quantified by a counter. The IC_{50} values are calculated as the concentration of the test compound at which the c-Jun phosphorylation is reduced to 50% of the control value. Preferred compounds of the present invention have an IC_{50} value ranging 0.01 - 10 μ M in this assay. To this end, a
- 25 preferred compound of this invention is Compound 1, which has an IC_{50} according to this assay of 0.11 μ M for JNK1 and JNK2, and 0.15 μ M for JNK3.

Selectivity For JNK

Compound 1 was also assayed for its inhibitory activity against the following protein kinases by techniques known to those skilled in this field (*see, e.g., Protein Phosphorylation*, Sefton & Hunter, Eds., Academic Press, pp. 97-367, 1998):

<u>Enzyme</u>	<u>IC₅₀</u>
p38-2	>30,000 nM
ERK1	>30,000 nM
MEKK1	>30,000 nM
IKK1	>30,000 nM
IKK2	>30,000 nM
PKA	>30,000 nM
PKC	>10,000 nM
EGF-TK	>10,000 nM

5

Jurkat T-cell IL-2 Production Assay

Jurkat T cells (clone E6-1) are purchased from the American Tissue Culture Collection and maintained in growth media consisting of RPMI 1640 medium containing 2 mM L-glutamine (Mediatech), with 10% fetal bovine serum (Hyclone) and penicillin/streptomycin. All cells are cultured at 37°C in 95% air and 5% CO₂. Cells are plated at a density of 0.2×10^6 cells per well in 200 µL of media. Compound stock (20 mM) is diluted in growth media and added to each well as a 10x concentrated solution in a volume of 25 µL, mixed, and allowed to pre-incubate with cells for 30 minutes. The compound vehicle (dimethylsulfoxide) is maintained at a final concentration of 0.5% in all samples. After 30 minutes the cells are activated with PMA (phorbol myristate acetate; final concentration 50 ng/mL) and PHA (phytohemagglutinin; final concentration 2 µg/mL). PMA and PHA are added as a 10x concentrated solution made up in growth media and added in a volume of 25 µL per well. Cell plates are cultured for 10 hours. Cells are pelleted by centrifugation and the media removed and stored at -20 °C. Media aliquots are analyzed by sandwich ELISA for the presence of IL-2 as per the manufacturers instructions (Endogen). The IC₅₀

20

values are calculated as the concentration of the test compound at which the IL-2 production was reduced to 50% of the control value. Preferred compounds of the present invention have an IC_{50} value ranging 0.1 - 30 μ M in this assay. Figure 1 presents the dose dependent inhibition of IL-2 in Jarkat T-Cells by Compound 1
5 according to this procedure, with a resulting IC_{50} of 5 μ M

Mouse *in vivo* LPS-Induced TNF- α Production Assay

Non-fasted mice are acclimatized for at least 7 days. Groups of 4 to 6 female BALB/c or CD-1 mice (8-10 weeks of age from Charles River laboratories) are
10 pretreated with test compound, either by intravenous injection or by oral gavage 15 – 180 minutes prior to the injection of 0.5 mg/kg Bacto LPS from *E. coli* 055:B5 (Difco Labs). Ninety minutes after LPS challenge, a terminal bleed is performed via abdominal vena cava and blood is allowed to clot at room temperature for 30 minutes in Microtainer serum separator tubes. After separation by centrifugation, the serum is
15 stored frozen at -80°C. ELIZA is performed on thawed, diluted samples (1:10 to 1:20) using a Mouse TNF-alpha kit (Biosource International). The ED_{50} values are calculated as the dose of the test compound at which the TNF- α production is reduced to 50% of the control value. Preferred compounds of the present invention have an ED_{50} value ranging 1 - 30 mg/kg in this assay. Figure 2 illustrates the results of this experiment
20 utilizing Compound 1 administered by intravenous injection (I.V.) at 15 and 30 mg/kg, as well as by per os (P.O.) at 7.5, 15 and 30 mg/kg. Vehicle alone (PEG-400, propylene glycol, cremophor EL, and ethanol in normal saline, "PPCES") and dexamethasone-21 acetate ("DEX") (1 mg/kg P.O.) were run as controls (n = 6, * = p 0.01). Compound 1 was administered 15 minutes pre-LPS challenge, and bleed occurred 90 minutes post
25 LPS.

Inhibition of Leukocyte Recruitment in Rat Inflamed Lung

Aerosol administration of ovalbumin in Brown Norway Rats previously sensitized by injection of ovalbumin (OA) results in an allergic airway inflammation
30 marked by the generation of an eosinophil- and T-lymphocyte-rich leukocytic

infiltration in the lungs (see Richards et al., *Am. J. Physiol.* 271:2 Pt 1, L267-76, 1996). Compound 1 was administered by subcutaneous injection at a dose of 30 mg/kg, b.i.d. for 3 days prior to ovalbumin challenge by aerosol. Cells counts were obtained from samples of broncho-alveolar lavage, the results of which are illustrated in Figure 3 (V =
5 PPCES vehicle).

Rat *In Vivo* Adjuvant Arthritis

Male Lewis rats were immunized with complete Freund's adjuvant on day 0 to induce an aggressive arthritis characterized by joint destruction and paw
10 swelling. Compound 1 was administered subcutaneously once daily from day 8 to day 20. Paw swelling was determined by water displacement plethysmometry (see Figure 4A; * = $p < 0.01$). Radiographs were obtained of the right hind paw to assess bone changes using a semi-quantitative scoring system: demineralization (0-2+), calcaneal
erosion (0-1+), and heterotopic bone formation (0-1+), with a maximum possible score
15 = 6 (see Figure 4B). Activation of AP-1 (see Figure 4C) was determined by DNA binding activity in an electrophoretic mobility shift assay (EMSA) (Ausubel et al., *Short Protocols in Molecular Biology*, Second Edition, John Wiley & Sons Publisher, New York, 1992). Matrix metalloproteinase-13 expression (see Figure 4D) was measured by
Northern blot analysis of MMP-13 mRNA (Ausubel et al., *supra*) (see also Winter et al.,
20 *Arthritis and Rheumatism* 9(3):394-404, 1966; Weichman et al., *Pharmacological Methods in the Control of Inflammation*, Chang and Lewis Eds., Alan R. Liss, Inc., Publ., New York, 1989).

Kainic Acid-Induced Seizure Response


25 Compound 1 was administered to male CD rats at 10 mg/kg intravenously through a tail vein catheter. This was followed immediately by a 30 mg/kg subcutaneous injection. Vehicle controls received the same injection volumes of the PPCES vehicle alone. Thirty minutes later, animals were given a 1- mg/kg i.p. injection of kainic acid in normal saline solution. This dose of kainic acid has been
30 previously reported to induce a seizure syndrome in rats (Maj et al., *Eur. J. Pharm.*

359:27-32, 1992). Seizure behavior was monitored for 4 hours following kainic acid injection. As presented in Figure 5, behaviors were assessed based on the following cumulative scoring system: 1 pt. = arrest of motion; 2 pts. = myoclonic jerks of the head and neck (moderate); 3 pts. = unilateral or bilateral forelimb clonic activity; 4 pts.
5 = whole body clonus; 5 pts. = clonic-tonic seizures; 6 pts. = status epilepticus (*see also* Mathis and Ungerer, *Exp. Brain Res.* 88:277-282, 1992; Rong et al., *Proc. Natl. Acad. Sci. USA* 96:9897-9902, 1999; Yang et al., *Nature* 389:865-870, 1997)

It will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications
10 may be made without departing from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS


1. A compound having the structure:



or a pharmaceutically acceptable salt thereof,

wherein

R_1 and R_2 are optional substituents that are the same or different and independently represent alkyl, halogen, nitro, trifluoromethyl, sulfonyl, carboxyl, alkoxycarbonyl, alkoxy, aryl, aryloxy, arylalkyloxy, arylalkyl, cycloalkylalkyloxy, cycloalkyloxy, alkoxyalkyl, alkoxyalkoxy, aminoalkoxy, mono- or di-alkylaminoalkoxy, or a group represented by formula (a), (b), (c) or (d):




R_3 and R_4 taken together represent alkylidene or a heteroatom-containing alkylidene, or R_3 and R_4 are the same or different and independently represent hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, aryloxyalkyl, alkoxyalkyl, alkoxyamino, or alkoxy(mono- or di-alkylamino); and


R_1 represents hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, alkoxy, amino, mono- or di-alkylamino, arylamino, arylalkylamino, cycloalkylamino, or cycloalkylalkylamino;

and with the proviso that at least R_1 or R_2 is present.


2. The compound of claim 1 wherein R_1 or R_2 is present, and having one of the following structures:




3. The compound of claim 1 wherein both R_1 and R_2 are present, and having one of the following structures:




4. The compound of claim 2 wherein R_1 and R_2 are:




5. The compound of claim 2 wherein R_1 and R_2 are:



6. The compound of claim 2 wherein R_1 and R_2 are:




7. The compound of claim 2 wherein R_1 and R_2 are:



8. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier.


9. A method for treating a condition responsive to JNK inhibition, comprising administering to a patient in need thereof and effective amount of a compound having the structure:



or a pharmaceutically acceptable salt thereof,

wherein

R_1 and R_2 are optional substituents that are the same or different and independently represent alkyl, halogen, nitro, trifluoromethyl, sulfonyl, carboxyl, alkoxycarbonyl, alkoxy, aryl, aryloxy, arylalkyloxy, arylalkyl, cycloalkylalkyloxy, cycloalkyloxy, alkoxyalkyl, alkoxyalkoxy, aminoalkoxy, mono- or di-alkylaminoalkoxy, or a group represented by formula (a), (b), (c) or (d):




R_3 and R_4 taken together represent alkylidene or a heteroatom-containing alkylidene, or R_3 and R_4 are the same or different and independently represent hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, aryloxyalkyl, alkoxyalkyl, alkoxyamino, or alkoxy(mono- or di-alkylamino); and



R, represents hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, cycloalkylalkyl, alkoxy, amino, mono- or di-alkylamino, arylamino, arylalkylamino, cycloalkylamino, or cycloalkylalkylamino.

10. The method of claim 9 wherein the condition is cancer.
11. The method of claim 9 wherein the condition is rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gout, asthma, bronchitis, cystic fibrosis, inflammatory bowel disease, irritable bowel syndrome, mucous colitis, ulcerative colitis, Crohn's disease, gastritis, esophagitis, hepatitis, multiple sclerosis, endotoxin shock, psoriasis, eczema, or dermatitis.
12. The method of claim 9 wherein the condition is atherosclerosis, restenosis following angioplasty, left ventricular hypertrophy, or myocardial infarction.
13. The method of claim 9 wherein the condition is stroke or ischemic damage to the heart, kidney, liver, or brain.
14. The method of claim 9 wherein the condition is transplant rejection.
15. The method of claim 9 wherein the condition is a central or peripheral neurological degenerative disorder.
16. The method of claim 15 wherein the central or peripheral neurological degenerative disorder is epilepsy, Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, a peripheral neuropathy, or spinal cord damage.




17. The method of claim 9 wherein R_1 and R_2 are not present, and having the following structure:



18. The method of claim 9 wherein R_1 or R_2 is present, and having one of the following structures:




19. The method of claim 9 wherein both R_1 and R_2 are present, and having one of the following structures:




20. The method of claim 18 wherein R_1 and R_2 are:


58




21. The method of claim 18 wherein R_1 and R_2 are:



22. The method of claim 18 wherein R_1 and R_2 are:




23. The method of claim 18 wherein R_1 and R_2 are:



24. A composition comprising a compound having the structure:

59



or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

Figure 1




Figure 2




Figure 3




Figure 4A





Figure 4B



Compound 1				Vehicle			
Control							
1	2	3	4	5	6	7	8

AP-1




Figure 4C

Vehicle				Compound 1			
1	2	3	4	5	6	7	8

MMP-13



Figure 4D

Figure 5

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.
PCT/US 00/22727

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D231/54 C07D401/04 C07D413/04 C07D401/12 C07D409/12
A61K31/415

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; HARTLEY, JOHN A. ET AL: "Characteristics of the interaction of anthrapyrazole anticancer agents with deoxyribonucleic acids: structural requirements for DNA binding, intercalation, and photosensitization" retrieved from STN Database accession no. 109:216 XP002152983 R3=NH(CH2)2NH2 abstract & MOL. PHARMACOL. (1988), 33(3), 265-71 ,</p> <p style="text-align: center;">--- -/--</p>	1,8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- 'A' document defining the general state of the art which is not considered to be of particular relevance
- 'E' earlier document but published on or after the international filing date
- 'L' document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- 'O' document referring to an oral disclosure, use, exhibition or other means
- 'P' document published prior to the international filing date but later than the priority date claimed

- 'T' later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- 'X' document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- '&' document member of the same patent family

Date of the actual completion of the international search

17 November 2000

Date of mailing of the international search report

04/12/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
Fax (+31-70) 340-3018

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 00/22727

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SHOWALTER, H. D. HOLLIS ET AL: "Anthrapyrazole anticancer agents. Synthesis and structure-activity relationships against murine leukemias" retrieved from STN Database accession no. 106:32913 XP002152984 RN 64598-58-9,91440-29-8,104739-78-8 abstract & J. MED. CHEM. (1987), 30(1), 121-31 ,</p>	1,8
X	<p>--- DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; IVANOVA, T. ET AL: "XPS investigation of electronic structure of pyrazolanthrone and its derivatives" retrieved from STN Database accession no. 127:176108 XP002152985 RNs 65846-98-2,70173-75-0,109185-28-6,163359-0 6-6 & POVERKHNOST (1997), (4-5), 193-201 ,</p>	1
X	<p>--- DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; GVON, HAN IR ET AL: "Amino-imino tautomerism and intramolecular cyclization of 4,9-diamino-1,10-anthraquinone-1-tosylimin es" retrieved from STN Database accession no. 122:160236 XP002152986 RNs 159790-09-7,109963-93-1 & DOKL. AKAD. NAUK (1994), 334(4), 465-8 ,</p> <p>--- -/-</p>	1

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.

PCT/US 00/22727

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SOKOLYUK, N. T. ET AL: "Synthesis and photochemical properties of peri-phenoxy derivatives of 6H-anthra[1,9-cd]-6-pyrazolone (pyrazole anthrone)" retrieved from STN Database accession no. 120:30709 XP002152987 Rns 56150-01-7, 64598-58-9, 151803-92-8, 151803-9 3-9 & ZH. ORG. KHIM. (1992), 28(10), 2193-200</p>	1
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; AMES, JAMES R. ET AL: "An integrated concept of amebicidal action: electron transfer and oxy radicals" retrieved from STN Database accession no. 107:190407 XP002152988 RN 109963-93-1 & FREE RADICAL BIOL. MED. (1987), 3(2), 85-96 ,</p>	1,8
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SINGH, M. R. ET AL: "Reactions of 2,2'-ethylenebis(anthrapyrazolone)" retrieved from STN Database accession no. 89:112272 XP002152989 Rns 67039-07-0, 67039-08-1, 67072-00-8, & INDIAN J. CHEM., SECT. B (1978), 16B(2), 100-2 ,</p>	1

-/--

INTERNATIONAL SEARCH REPORT

Intern. Appl. Application No

PCT/US 00/22727

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; GALUSHKO, A. M. ET AL: "Derivatives of pyrazoloanthrone. I. Reactivity of 3-aminopyrazoloanthrone" retrieved from STN Database accession no. 87:184421 XP002152990 RNs 64573-16-6, 64573-17-7, 64598-58-9 & KHIM. GETEROTSIKL. SOEDIN. (1977), (7), 956-61 ,</p>	1
X	<p>FR 2 401 915 A (CIBA GEIGY AG) 30 March 1979 (1979-03-30) compounds (3), (4), (5)</p>	1
X	<p>FR 2 336 708 A (HOECHST AG) 22 July 1977 (1977-07-22) example 30</p>	1
X	<p>FR 2 167 626 A (BASF AG) 24 August 1973 (1973-08-24) example 19</p>	1
X	<p>FR 2 024 807 A (CIBA GEIGY) 4 September 1970 (1970-09-04) example 14 (RN 41313-85-3)</p>	1
X	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; ARIENT, JOSEF: "5,8-Dimethylpyrazoloanthrone" retrieved from STN Database accession no. 79:6796 XP002152991 RN 40537-86-8 & CS 146 895 1 (ARIENT, JOSEF) 15 January 1973 (1973-01-15)</p>	1
A	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; JUDSON, IAN R.: "The anthrapyrazoles: a new class of compounds with clinical activity in breast cancer" retrieved from STN Database accession no. 118:139035 CA XP002152992 abstract & SEMIN. ONCOL. (1992), 19(6), 687-94 , 1992,</p>	1-24

-/-

INTERNATIONAL SEARCH REPORT

Inter. Appl. No.
PCT/US 00/22727

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SHOWALTER, H. D. HOLLIS ET AL: "5-(Aminoalkyl)amino!-substituted anthra'1,9-cd!pyrazol-6(2H)-ones as novel anticancer agents. Synthesis and biological evaluation" retrieved from STN Database accession no. 100:85627 CA XP002152993 abstract & J. MED. CHEM. (1984), 27(3), 253-5 , 1984,</p>	1-24

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Appl. Application No

PCT/US 00/22727

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2401915 A	30-03-1979	CH 633525 A	15-12-1982
		BR 7805710 A	17-04-1979
		CA 1109065 A	15-09-1981
		DE 2838090 A	15-03-1979
		ES 473018 A	16-03-1979
		GB 2003477 A,B	14-03-1979
		JP 54052076 A	24-04-1979
		US 4198518 A	15-04-1980
		BE 870138 A	01-03-1979
		BE 870139 A	01-03-1979
		BR 7805716 A	17-04-1979
		CA 1108609 A	08-09-1981
		DE 2838091 A	15-03-1979
		ES 473017 A	16-03-1979
		FR 2401908 A	30-03-1979
		GB 2003476 A,B	14-03-1979
		JP 54048746 A	17-04-1979
		US 4202827 A	13-05-1980
FR 2336708 A	22-07-1977	DE 2558813 A	07-07-1977
		BE 849869 A	24-06-1977
		CA 1058943 A	24-07-1979
		GB 1576217 A	01-10-1980
		JP 1258089 C	29-03-1985
		JP 52082415 A	09-07-1977
		JP 59033893 B	18-08-1984
FR 2167626 A	24-08-1973	NL 7614408 A,B,	29-06-1977
		DE 2200127 A	12-07-1973
		CH 575364 A	14-05-1976
		GB 1404969 A	03-09-1975
		IT 974374 B	20-06-1974
FR 2024807 A	04-09-1970	CH 538527 A	15-08-1973
		DE 1954453 A	27-08-1970
		ES 373225 A	16-04-1972
		GB 1293557 A	18-10-1972
		GB 1293558 A	18-10-1972
CS 146895	1	NONE	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.